

In vitro maturation: a committee opinion

The Practice Committees of the American Society for Reproductive Medicine and the Society for Assisted Reproductive Technology

American Society for Reproductive Medicine, Birmingham, Alabama

The initial results of in vitro maturation suggest the potential for clinical application. However, at this time in vitro maturation should be performed only as an experimental procedure evaluating both efficacy and safety in carefully selected patients. (Fertil Steril® 2013;99:663–6. ©2013 by American Society for Reproductive Medicine.)

Earn online CME credit related to this document at www.asrm.org/elearn

Discuss: You can discuss this article with its authors and with other ASRM members at <http://fertstertforum.com/goldsteinj-in-vitro-maturation-committee-opinion/>



Use your smartphone to scan this QR code and connect to the discussion forum for this article now.*

* Download a free QR code scanner by searching for "QR scanner" in your smartphone's app store or app marketplace.

In vitro fertilization usually involves ovarian stimulation with exogenous gonadotropins to maintain the development of gonadotropin-sensitive follicles and inhibit the atresia of the non-dominant follicles (1, 2). The term in vitro maturation (IVM) refers to the maturation in culture of immature oocytes after their recovery from follicles that may or may not have been exposed to exogenous follicle-stimulating hormone (FSH) but were not exposed to either exogenous luteinizing hormone (LH) or human chorionic gonadotropin (hCG) prior to retrieval to induce meiotic resumption. During IVM, such immature oocytes progress from prophase I stage (i.e., from the germinal vesicle [GV]) through meiosis I to reach metaphase II (MII). The use of exogenous LH or hCG prior to retrieval ("follicular priming") has been introduced into clinical applications in order to increase the likelihood of obtaining usable oocytes (3–5). While this practice is not IVM in the strictest sense, outcomes with these methods will be included in this document.

The human oocyte reaches its full size (approximately 100 μm to 120 μm) at the antral stage, during which time the follicular diameter is only a fraction of its final ovulatory diameter. The ability of an oocyte to resume and complete meiosis is linked closely to follicular diameter (6). In humans, the kinetics of oocyte maturation have been established with oocytes obtained from oophorectomy specimens from nonmalignant gynecological disorders and then cultured to undergo IVM (Fig. 1) (3, 4, 7, 8).

It should be noted that even if an immature oocyte progresses to the MII stage, complete oocyte competence is not necessarily obtained. In many cases, oocyte maturation in vitro is accomplished morphologically but without acquiring developmental competence following in vitro culture. In order for a mature oocyte to undergo successful fertilization and subsequent development, synchronization of nuclear and cytoplasmic maturation must occur. Nuclear maturation consists of germinal vesicle breakdown induced by the LH surge followed by resumption of

meiosis and extrusion of the first polar body (MII). Cytoplasmic maturation is more difficult to assess microscopically but refers to an accumulation of factors that prepare the cytoplasm for fertilization and embryonic development (9). Epigenetic processes are a component of nuclear and cytoplasmic oocyte maturation, influencing development after fertilization (10, 11). The potential for abnormal methylation of maternally expressed genes needs careful evaluation in oocytes matured in vitro.

POTENTIAL APPLICATIONS OF IVM

Potential indications for IVM include patients at risk for ovarian hyperstimulation syndrome (OHSS), those with limited time for ovarian stimulation, and those with a contraindication to sustained elevations of estradiol (E2).

Since IVM either omits gonadotropin stimulation or uses a short course of gonadotropins, there is a reduction in the number of days of follicular monitoring, utilization of gonadotropins, and the risk of OHSS. Women with polycystic ovary syndrome (PCOS) or PCO-like ovaries may benefit from IVM. These women have the highest number of antral follicles for potential retrieval and are at highest risk of developing OHSS under traditional stimulation protocols.


Received December 17, 2012; accepted December 19, 2012; published online February 1, 2013.

No reprints will be available.

Correspondence: Practice Committee, American Society for Reproductive Medicine, 1209 Montgomery Hwy., Birmingham, Alabama 35216 (E-mail: ASRM@asrm.org).

Fertility and Sterility® Vol. 99, No. 3, March 1, 2013 0015-0282/\$36.00

Copyright ©2013 American Society for Reproductive Medicine, Published by Elsevier Inc. <http://dx.doi.org/10.1016/j.fertnstert.2012.12.031>

FIGURE 1

IVM can be used in patients who need to initiate gonadotoxic therapies without adequate time to undergo ovarian stimulation with retrieval of oocytes matured in vivo. Patients with hormonally sensitive tumors or other contraindications to sustained elevations of E2 also may consider using IVM in an effort to avoid the increased E2 concentrations achieved with ovarian stimulation.

The applicability of these indications may be limited by alternative strategies to minimize OHSS risk and, in the case of those with limited time, the delay incurred by coordinating treatment with the onset of the menstrual cycle, even in the absence of ovarian stimulation.

IVM IN CLINICAL PRACTICE

Patient Preparation

No randomized, controlled trials currently exist comparing IVM with IVF; however, there have been a number of clinical studies using in vitro matured human oocytes in PCO-like and PCOS patients (12–20) and in ovulatory women (15, 20–24). Several trials also have evaluated the utility of follicular priming versus the retrieval of oocytes from unstimulated antral follicles (12, 18–21, 25).

Three methods of follicular priming have been reported. One method is to prime oocytes with FSH for 3–6 days, followed by retrieval on cycle day 9–10; this is the strictest definition of IVM. Other approaches utilize hCG priming (and include single injections of 10,000 IU of hCG to “prime” intermediate-sized follicles 36 hours prior to oocyte retrieval) and the use of FSH stimulation in addition to hCG. Priming with hCG or FSH gives mixed results; however, priming with either agent appears to benefit PCOS patients in terms of implantation and pregnancy rates (14, 17, 19) over no priming at all (Table 1). The combination of gonadotropin and hCG priming compared to single-agent follicular priming has been studied with disparate results. While no difference was seen in one trial (18), a large randomized trial showed

that the combination of FSH and hCG improved the percentage of oocytes that matured in vitro as well as implantation rates when compared to no priming or hCG alone. The clinical pregnancy rates were highest in the combination group (26%) compared to 11%, 5%, and 13% in the not primed, hCG-primed, or FSH-primed groups, respectively. The implantation rate of 16.4% was also significantly higher than in the no priming (9.2%) or the hCG-primed (4.0%) but not the FSH-primed group (10.6%) (25).

Interpretation of these studies is impaired by inconsistent reporting of the developmental stage of the oocytes at retrieval and the status of the cumulus at assessment and fertilization.

Oocyte Retrieval and Culture

There are no established criteria to identify the ideal timing or method for oocyte retrieval, with most studies using a lead-follicular diameter of up to 10 mm (17, 24, 26). Lead-follicle diameters greater than 13 mm have been associated with reduced numbers of oocytes collected and matured (27), possibly related to subsequent atresia of the non-dominant follicles from withdrawal of endogenous FSH support. Some protocols require an endometrial thickness of >5 mm while others do not include endometrial thickness as a criterion.

The aspiration technique also differs for immature oocytes compared to traditional IVF. Double lumen needles to flush follicles have been described but do not appear to be necessary to retrieve immature oocytes. The optimal aspiration pressure and needle design have yet to be determined with negative pressures ranging between 80 mm Hg and 300 mm Hg and needles ranging from 16 to 20 gauge described (24, 28). Extremely high aspiration pressure has been shown to negatively impact development (29).

There is no consensus on which formulation is best suited for the purpose of in vitro oocyte culture. Media are frequently supplemented with pyruvic acid along with essential and non-essential amino acids, such as tissue culture media (TCM).

Fertilization

Because of potential changes in the characteristics of the oocytes and zona pellucidae from culture of immature oocytes, intracytoplasmic sperm injection (ICSI) has been advocated as the preferred method for fertilization. While fertilization rates appear to be increased utilizing ICSI for IVM oocytes, developmental competence may be impaired as demonstrated in one comparative trial (20). Fertilization rates of matured oocytes in patients who did not receive gonadotropins were only 37.7% (229/608 matured oocytes) with conventional IVF compared to 69.3% (318/459 mature oocytes) when ICSI was used as the insemination technique. Despite lower fertilization results, the implantation rate was significantly higher in embryos derived from oocytes fertilized with conventional IVF compared to ICSI (24.2% vs. 14.8%; $P < .05$) as well as clinical pregnancy rates per embryo transfer procedure (34.5% vs. 20.0%; $P < .05$). Comparative trials comparing IVF to ICSI for fertilization of in vitro matured oocytes need to be conducted.

TABLE 1

In vitro maturation in PCOS and PCO-like patients.

Reference	N	Priming	Mean no. oocytes	Mean no. ET	PR/ET, %	IR, %	MC, %
13	203	None	15.5	5.0	21.9	5.5	37
14	13	hCG	7.8	2.8	38.5	16.6	40
	11	None	7.4	2.5	27.3	14.8	0
16	107	hCG	10.3	2.5	21.5	9.5	26
19	12	None	6.8	1.7	0	0	0
	24	FSH	6.5	1.8	33.0	21.6	63
17	45	hCG	11.4	2.5	22.5	10.9	40
18	35	FSH/hCG	21.9	3.8	31.4	9.7	13
	33	hCG	23.1	3.8	36.4	11.3	?

Note: ET = embryos transferred; PR/ET = pregnancies per embryo transfer procedure; IR = implantation rate (sacs/embryos transferred); MC = miscarriage rate.

Practice Committee. In vitro maturation. *Fertil Steril* 2013.

Clinical Outcome

Two large series have been reported (30). In the first, a total of 3,079 oocytes collected from PCOS patients following hCG priming, 78.8% of immature oocytes reached maturity and 69.2% successfully fertilized with ICSI. In the second, 73.2% of 6,860 oocytes matured in vitro following hCG priming with a fertilization rate of 79.0%. Although clinical pregnancy rates in these and other IVM series approach that of traditional IVF (23% to 34%), this is secondary to the increased number of embryos transferred (Table 1). As a consequence, implantation rates are a more reliable indicator of IVM efficiency and range between 5.5% and 21.6%, lower than expected for women of comparable age undergoing conventional IVF (Table 1). In the Society for Assisted Reproductive Technology (SART) report from cycles performed in 2010 in young women (<35 years of age), implantation rates of 36.9% and a mean number of embryos transferred of 2.0 yields a live birth rate per started cycle of 41.7% (31).

The developmental outcome of children conceived with IVM has been studied in small numbers and thus far are reported to be no different than children born through traditional IVF or ICSI (32, 33). However, the small number of children conceived through IVM limits the accuracy of malformation and anomaly rates, and developmental outcomes cannot be assessed accurately.

SUMMARY

- Although there are no RCT comparisons, available data indicate that IVM as described in this document results in lower-than-expected success rates.
- In many cases, nuclear maturation in vitro is evident; however, there is a failure to acquire developmental competence following in vitro culture secondary to cytoplasmic competence.
- The implantation and pregnancy rates of IVM are less than expected with ovarian stimulation and retrieval of in vivo matured oocytes.
- Because only a small number of children have been conceived with IVM, information on the safety of IVM with regard to malformation and developmental outcomes cannot be assessed accurately.

- Priming with hCG and/or FSH appears to improve implantation and pregnancy rates compared to no priming.

CONCLUSIONS

- Candidates for this technology include those at risk for OHSS, women with PCOS or PCO-like ovaries, women with estrogen-sensitive cancers, or those with limited time prior to initiating potentially gonadotoxic treatments for fertility preservation.
- ICSI does not appear to be necessary to achieve fertilization in oocytes matured in vitro.
- The initial results of in vitro maturation suggest the potential for clinical application. However, at this time, patients must be made aware that the implantation and pregnancy rates are significantly lower than with standard IVF, limiting more universal utilization.
- In vitro maturation should only be performed as an experimental procedure (34) in specialized centers for carefully selected patients evaluating both efficacy and safety. Informed consent must include information regarding pregnancy rates of IVM in comparison to conventional ART and alternative options, if any.

Acknowledgments: This report was developed under the direction of the Practice Committees of the American Society for Reproductive Medicine and the Society for Assisted Reproductive Technology as a service to its members and other practicing clinicians. Although this document reflects appropriate management of a problem encountered in the practice of reproductive medicine, it is not intended to be the only approved standard of practice or to dictate an exclusive course of treatment. Other plans of management may be appropriate, taking into account the needs of the individual patient, available resources, and institutional or clinical practice limitations. The Practice Committees and the Board of Directors of the American Society for Reproductive Medicine and the Society for Assisted Reproductive Technology have approved this report.

The following members of the ASRM Practice Committee participated in the development of this document. All committee members disclosed commercial and financial relationships with manufacturers or distributors of goods or services

used to treat patients. Members of the Committee who were found to have conflicts of interest based on the relationships disclosed did not participate in the discussion or development of this document.

Samantha Pfeifer, M.D.; Marc Fritz, M.D.; Jeffrey Goldberg, M.D.; G. David Adamson, M.D.; R. Dale McClure, M.D.; Roger Lobo, M.D.; Michael Thomas, M.D.; Eric Widra, M.D.; Mark Licht, M.D.; John Collins, M.D.; Marcelle Cedars, M.D.; Catherine Racowsky, Ph.D.; Michael Vernon, M.D.; Owen Davis, M.D.; Kurt Barnhardt, M.D., M.S.C.E.; Glenn Schattman, M.D.; Clarisa Gracia, M.D., M.S.C.E.; William Catherino, M.D., Ph.D.; Kim Thornton, M.D.; Robert Rebar, M.D.; Andrew La Barbera, Ph.D.

REFERENCES

1. Cha KY, Chian RC. Maturation in vitro of immature human oocytes for clinical use. *Hum Reprod Update* 1998;4:103–20.
2. Russell JB, Knezevich KM, Fabian KF, Dickson JA. Unstimulated immature oocyte retrieval: early versus midfollicular endometrial priming. *Fertil Steril* 1997;67:616–620.
3. Anderiesz C, Trouson AO. The effect of testosterone on the maturation and developmental capacity of murine oocytes in vitro. *Hum Reprod* 1995;10:2377–81.
4. Baker SJ, Spears N. The role of intra-ovarian interactions in the regulation of follicular dominance. *Hum Reprod Update* 1999;5:153–65.
5. Cha KY, Koo JJ, Ko JJ, Choi DH, Han SY, Yoon TK. Pregnancy after in vitro fertilization of human follicular oocytes collected from non-stimulated cycles, their culture in vitro and their transfer in a donor-oocyte program. *Fertil Steril* 1991;55:109–13.
6. Gilchrist RB, Nayudu PL, Nowshari MA, Hodges JK. Meiotic competence of marmoset monkey oocytes is related to follicle size and oocyte-somatic cell associations. *Biol Reprod* 1995;52:1234–43.
7. Edwards RG. Maturation in vitro of human ovarian oocytes. *Lancet* 1965;286:926–9.
8. Gunnala S, Kaufman ML, Racowsky C. Kinetics of meiotic maturation in cultured human oocytes retrieved following oophorectomy. *Assis Reprod Technol Androl* 1993;4:1–14.
9. Moor RM, Dai Y, Lee C, Fulka J Jr. Oocyte maturation and embryonic failure. *Hum Reprod Update* 1998;4:223–36.
10. Borghol N, Lornage J, Blachere T, Garret SA, Lefevre A. Epigenetic status of the H19 locus in human oocytes following in vitro maturation. *Genomics* 2006;87:417–26.
11. Bromfield J, Messamore W, Albertini DF. Epigenetic regulation during mammalian oogenesis. *Reprod Fert and Develop* 2008;20:74–80.
12. Cha KY, Han SY, Chung HM, Choi DH, Lim JM, Lee WS, et al. Pregnancies and deliveries after in vitro maturation culture followed by *in vitro* fertilization and embryo transfer without stimulation in women with polycystic ovary syndrome. *Fertil Steril* 2000;73:978–83.
13. Cha KY, Chung HM, Lee DR, Kwon H, Chung MK, Park LS, et al. Obstetric outcome of patients with polycystic ovary syndrome treated by in vitro maturation and *in vitro* fertilization-embryo transfer. *Fertil Steril* 2005;83:1461–5.
14. Chian RC, Buckett WM, Tulandi T, Tan SL. Prospective randomized study of human chorionic gonadotropin priming of women with polycystic ovarian syndrome. *Hum Reprod* 2000;15:165–70.
15. Child TJ, Abdul-Jalil AK, Gulekli B, Tan SL. In vitro maturation and fertilization of oocytes from unstimulated normal ovaries, polycystic ovaries, and women with polycystic ovarian syndrome. *Fertil Steril* 2001;76:936–42.
16. Child TJ, Phillips SJ, Abdul-Jalil AK, Gulekli B, Tan SL. A comparison of in vitro maturation and in vitro fertilization for women with polycystic ovaries. *Obstet Gynecol* 2002;100:665–70.
17. Le Du A, Kadoch IJ, Bourcigaux N, Doumerc S, Bourruer MC, Chevalier N, et al. In vitro maturation for the treatment of infertility associated with polycystic ovarian syndrome: the French experience. *Hum Reprod* 2005;20:420–4.
18. Lin YH, Hwang JL, Huang LW, Mu SC, Seow KM, Chung J, et al. Combination of FSH priming and hCG priming for in vitro maturation of human oocytes. *Hum Reprod* 2003;18:1632–6.
19. Mikkelsen AL, Lindenberg S. Benefit of hCG priming of women with PCOS to the in vitro maturation procedure and the outcome: a randomized prospective study. *Reproduction* 2001;122:587–92.
20. Soderstrom-Anttila V, Makinen S, Tuuri T, Suikkari AM. Favourable pregnancy results with insemination of in vitro matured oocytes from unstimulated patients. *Hum Reprod* 2005;20:1534–40.
21. Mikkelsen AL, Smith SD, Lindenberg S. In-vitro maturation of human oocytes from regularly menstruating women may be successful without follicle stimulating hormone priming. *Hum Reprod* 1999;14:1847–51.
22. Mikkelsen AL, Smith S, Lindenberg S. Impact of oestradiol and inhibin A concentrations on pregnancy rate in in-vitro oocyte maturation. *Hum Reprod* 2000;15:1685–90.
23. Mikkelsen AL, Andersson AM, Skakkebaek NE, Lindenberg S. Basal concentrations of oestradiol may predict the outcome of in-vitro maturation in regularly menstruating women. *Hum Reprod* 2001;16:862–7.
24. Yoon HG, Yoon SH, Son WY, Lee SW, Park SP, Im KS, et al. Pregnancies resulting from in vitro matured oocytes collected from women with regular menstrual cycle. *J Assist Reprod Genet* 2001;18:325–9.
25. Fadini R, Del Canto MB, Renzini MM, Brambillasca F, Comi R, Fumagalli D, et al. Effect of different gonadotrophin priming on IVM of oocytes from women with normal ovaries: a prospective randomized study. *Reprod Biomed Online* 2009;19:343–51.
26. Cobo AC, Requena A, Neuspiller F, Aragones M, Mercader A, Navarro J, et al. Maturation in vitro of human oocytes from unstimulated cycles: selection of the optimal day for ovum retrieval based on follicular size. *Hum Reprod* 1999;14:1864–8.
27. Jurema M, Noguiera D. In vitro maturation of human oocytes for assisted reproduction. *Fertil Steril* 2006;86:1277–91.
28. Wynn P, Picton HM, Krapez JA, Rutherford AJ, Balen AH, Gosden RJ. Pre-treatment with follicle stimulating hormone promotes the numbers of human oocytes reaching metaphase II by in vitro maturation. *Hum Reprod* 1998;13:3132–8.
29. Hashimoto S, Fukuda A, Murata Y, Kikkawa M, Oku H, Kanaya H, et al. Effect of aspiration vacuum on the developmental competence of immature human oocytes retrieved using a 20-gauge needle. *Reprod Biomed Online* 2007;14:444–9.
30. Chian RC, Buckett WM, Tan SL. In-vitro maturation of human oocytes. *Reprod Biomed Online* 2004;8:148–66.
31. Centers for Disease Control and Prevention, American Society for Reproductive Medicine, Society for Assisted Reproductive Technology. 2010 Assisted Reproductive Technology success rates: national summary and fertility clinic reports. Atlanta: U.S. Department of Health and Human Services; 2010.
32. Söderström-Anttila V, Salokorpi T, Pihlaja M, Serenius-Sirve S, Suikkari AM. Obstetric and perinatal outcome and preliminary results of development of children born after in vitro maturation of oocytes. *Hum Reprod* 2006;21:1508–13.
33. Buckett WM, Chian RC, Holzer H, Dean N, Usher R, Tan SL. Obstetric outcomes and congenital abnormalities after in vitro maturation, in vitro fertilization, and intracytoplasmic sperm injection. *Obstet Gynecol* 2007;110:885–91.
34. Practice Committee of the American Society for Reproductive Medicine. Definition of “experimental procedures”. *Fertil Steril* 2009;92:1517.